

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Pyrophosphatase, Inorganic (E. coli)

Catalog Number: M0361S Concentration: 100 U/ml

Unit Definition: One unit is the amount of enzyme that will generate 1 µmol of

phosphate per minute from inorganic pyrophosphate under standard

reaction conditions.

Packaging Lot Number: 10065128
Expiration Date: 08/2021
Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl , 20 mM Tris-HCl (pH 8.0), 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol

Specification Version: PS-M0361S/L v2.0

Pyrophosphatase, Inorganic (E. coli) Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0361SVIAL	Pyrophosphatase, Inorganic (E. coli)	10052545	Pass	

Assay Name/Specification	Lot # 10065128
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 4 containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 0.5 units of Pyrophosphatase, Inorganic (E. coli) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Protein Purity Assay (SDS-PAGE) Pyrophosphatase, Inorganic (E. coli) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 4 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 0.5 units of Pyrophosphatase, Inorganic (E. coli) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 0.1 unit of Pyrophosphatase, Inorganic (E. coli) is incubated at	Pass



M0361S / Lot: 10065128

Page 1 of 2

Assay Name/Specification	Lot # 10065128
37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	
Phosphatase Activity (pNPP) A 100 µl reaction in NEBuffer 4 containing 10 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 1 unit Pyrophosphatase, Inorganic (E. coli) incubated for 1 hour at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass

This product has been tested and shown to be in compliance with all specifications.

Dongxian Yue

**Production Scientist** 

27 Sep 2019

Michael Tonello

Packaging Quality Control Inspector

22 Jan 2020

