

New England Biolabs Certificate of Analysis

Product Name: ProtoScript® II Reverse Transcriptase
Catalog Number: M0368X
Concentration: 200,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme that will incorporate 1 nmol of dTTP into an acid-insoluble form in 10 minutes at 37°C.
Packaging Lot Number: 10094714
Expiration Date: 03/2022
Storage Temperature: -20°C
Storage Conditions: 20 mM Tris-HCl, 100 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 0.01 % IGEPAL® CA-630, 50 % Glycerol, (pH 7.5 @ 25°C)
Specification Version: PS-M0368S/L/X v2.0

ProtoScript® II Reverse Transcriptase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0368L	ProtoScript® II Reverse Transcriptase	10087492	Pass

Assay Name/Specification	Lot # 10094714
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in ProtoScript® II Reverse Transcriptase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 200 units of ProtoScript® II Reverse Transcriptase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 200 units of ProtoScript® II Reverse Transcriptase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) ProtoScript® II Reverse Transcriptase is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 200 units of ProtoScript® II Reverse Transcriptase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from	Pass


Assay Name/Specification	Lot # 10094714
<p>purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p> <p>RNAse Activity Assay (4 Hour Digestion) A 10 μl reaction in ProtoScript[®] II Reverse Transcriptase Reaction Buffer containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μl of ProtoScript[®] II Reverse Transcriptase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p> <p>Endonuclease Activity (Nicking) A 50 μl reaction in ProtoScript[®] II Reverse Transcriptase Reaction Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 200 units of ProtoScript[®] II Reverse Transcriptase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<p style="text-align: center;">Pass</p> <p style="text-align: center;">Pass</p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Christie Vazquez
Production Scientist
29 Dec 2020



Michael Tonello
Packaging Quality Control Inspector
29 Dec 2020