240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: T3 RNA Polymerase

Catalog #: M0378S

Concentration: 50,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 1 nmol ATP into acid-insoluble material in a total reaction

volume of 50 µl in 1 hour at 37°C in 1X RNA Polymerase Reaction Buffer.

 Lot #:
 0011712

 Assay Date:
 12/2017

 Expiration Date:
 12/2019

 Storage Temp:
 -20°C

Storage Conditions: 50 mM Tris-HCl, 100 mM NaCl, 1 mM EDTA, 20 mM βME, 0.1% Triton®X-100, 50% Glycerol, (pH 7.9 @ 25°C)

Specification Version: PS-M0378S v1.0 Effective Date: 13 Jun 2018

Assay Name/Specification (minimum release criteria)	Lot #0011712
Endonuclease Activity (Nicking) - A 50 μl reaction in RNAPol Reaction Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 150 units of T3 RNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) - A 50 μ l reaction in RNAPol Reaction Buffer containing 1 μ g of a mixture of single and double-stranded [3 H] <i>E. coli</i> DNA and a minimum of 150 units of T3 RNA Polymerase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 μl reaction in RNAPol Reaction Buffer containing 1 μg of Lambda DNA and a minimum of 250 units of T3 RNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) - T3 RNA Polymerase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) - A 10 μ l reaction in RNAPol Reaction Buffer containing 40 ng of a 300 base single-stranded RNA and a minimum of 50 units of T3 RNA Polymerase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

Authorized by Derek Robinson 13 Jun 2018







Inspected by
Dongxian Yue
31 Dec 2017