

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	T3 RNA Polymerase
Catalog Number:	M0378S
Concentration:	50,000 U/mI
Unit Definition:	One unit is defined as the amount of enzyme that will incorporate 1 nmol ATP into acid-insoluble material in a total reaction volume of 50 μ l in 1 hour at 37°C in 1X RNA Polymerase Reaction Buffer.
Packaging Lot Number:	10110336
Expiration Date:	06/2023
Storage Temperature:	-20°C
Storage Conditions:	50 mM Tris-HCl, 100 mM NaCl, 1 mM EDTA, 20 mM βME, 0.1% Triton®X-100, 50% Glycerol, (pH 7.9 @ 25°C)
Specification Version:	PS-M0378S v1.0

T3 RNA Polymerase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0378SVIAL	T3 RNA Polymerase	10110335	Pass	
B9012SVIAL	RNAPol Reaction Buffer	10100016	Pass	

Assay Name/Specification	Lot # 10110336
RNase Activity (Extended Digestion) A 10 µl reaction in RNAPol Reaction Buffer containing 40 ng of a 300 base single-stranded RNA and a minimum of 50 units of T3 RNA Polymerase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Protein Purity Assay (SDS-PAGE) T3 RNA Polymerase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of Lambda DNA and a minimum of 250 units of T3 RNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μI reaction in RNAPol Reaction Buffer containing 1 μg of a mixture of single	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10110336
and double-stranded [³ H] E. coli DNA and a minimum of 150 units of T3 RNA Polymerase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Endonuclease Activity (Nicking) A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 150 units of T3 RNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Bhairavi Jani Production Scientist 03 Jun 2021

Michael 1. 11

Michael Tonello Packaging Quality Control Inspector 03 Jun 2021

