

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Cas9 Nuclease, S. pyogenes

Catalog Number: M0386S
Concentration: 1,000 nM
Packaging Lot Number: 10211544
Expiration Date: 09/2025
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl , 300 mM NaCl , 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0386S/L v1.0

Cas9 Nuclease, S. pyogenes Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0386SVIAL	Cas9 Nuclease, S. pyogenes	10204931	Pass	
B6003SVIAL	NEBuffer™ r3.1	10182164	Pass	

Assay Name/Specification	Lot # 10211544
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 3.1 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 1 pmol of Cas9 Nuclease, S. pyogenes incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 1 pmol of Cas9 Nuclease, S. pyogenes incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (Targeted Digestion) A 20 µl reaction in NEBuffer 3.1 containing 20 nM of 100 bp FAM and ROX-labeled double-stranded target DNA, 100 nM sgRNA, and 100 nM Cas9 Nuclease, S. pyogenes incubated for 1 hour at 37°C results in ≥90% targeted digestion of the substrate DNA as determined by capillary electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 1 pmol of Cas9 Nuclease, S. pyogenes incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel	Pass



M0386S / Lot: 10211544

Page 1 of 2

This product has been tested and shown to be in compliance with all specifications.

incubation for 16 hours, >90% of the substrate RNA remains intact as determined by

gel electrophoresis using fluorescent detection.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Jessica Cane **Production Scientist**

27 Sep 2023

Josh Hersey

Packaging Quality Control Inspector

03 Oct 2023



M0386S / Lot: 10211544

Page 2 of 2