

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: OneTaq® Quick-Load® DNA Polymerase

Catalog Number: M0509S Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 15

nmol of dNTP into acid insoluble material in 30 minutes at 75°C.

Packaging Lot Number: 10056355
Expiration Date: 07/2021
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.5 % Tween® 20,

0.5 % IGEPAL® CA-630, 50% Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0509S/L/X v1.0

OneTaq® Quick-Load® DNA Polymerase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0509SVIAL	OneTaq® Quick-Load® DNA Polymerase	10052145	Pass	
B9022SVIAL	OneTaq® Standard Reaction Buffer	10034011	Pass	
B0509AVIAL	OneTaq® Quick-Load® Reaction Buffer	10034017	Pass	

Assay Name/Specification	Lot # 10056355
PCR Amplification (5.0 kb Lambda DNA) A 25 μl reaction in OneTaq® Standard Reaction Buffer in the presence of 200 μM dNTPs and 0.2 μM primers containing 5 ng Lambda DNA with 0.625 units of OneTaq® Quick-Load® DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.	Pass
PCR Amplification (5.0 kb Lambda DNA) A 25 μl reaction in OneTaq® Quick-Load® Reaction Buffer in the presence of 200 μM dNTPs and 0.2 μM primers containing 5 ng Lambda DNA with 0.625 units of OneTaq® Quick-Load® DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of OneTaq® Quick-Load® DNA Polymerase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass



M0509S / Lot: 10056355

Page 1 of 2

Assay Name/Specification	Lot # 10056355
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction	
containing Lambda-HindIII DNA and a minimum of 5 units of OneTaq® Quick-Load® DNA	
Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

Christie Vazquez Production Scientist

vaistie Vazguez

26 Aug 2019

Michael Tonello

Packaging Quality Control Inspector

13 Jan 2020

