

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Quick CIP
Catalog Number:	M0525S
Concentration:	5,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme that hydrolyzes 1 μmol of p-Nitrophenyl Phosphate, PNPP in a total reaction volume of 1 ml in 1 minute at 37°C.
Lot Number:	10050886
Expiration Date:	06/2021
Storage Temperature:	-20°C
Storage Conditions:	25 mM Tris-HCl , 1 mM MgCl2 , 0.1 mM ZnCl2 , 50 % Glycerol, (pH 7.5 @ 25°C)
Specification Version:	PS-M0525S/L v1.0

Quick CIP Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0525SVIAL	Quick CIP	10047902	Pass	
B7204SVIAL	CutSmart® Buffer	10046087	Pass	

Assay Name/Specification	Lot # 10050886
RNase Activity (Extended Digestion) A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ l of Quick CIP is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using gel electrophoresis using fluorescent detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 μ I reaction in NEBuffer 4 containing 1 μ g of PhiX174-HaeIII DNA and a minimum of 50 units of Quick CIP incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 50 units of Quick CIP incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking)	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10050886
A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of Quick CIP incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

Differen Duquette

Production Scientist 09 Aug 2019

al-

An Minichiello Packaging Quality Control Inspector 20 Aug 2019

