


New England Biolabs Certificate of Analysis

Product Name: Exonuclease VIII, truncated
Catalog #: M0545S/L
Concentration: 10,000 units/ml
Unit Definition: One unit is defined as the amount of enzyme required to produce 1 nmol of acid-soluble deoxyribonucleotide from double-stranded DNA in a total reaction volume of 50 µl in 30 minutes at 37°C in 1X NEBuffer 4 with 0.15 mM sonicated duplex [³H]-DNA.
Lot #: 0031708
Assay Date: 08/2017
Expiration Date: 08/2019
Storage Temp: -20°C
Storage Conditions: 50 mM Tris-HCl, 100 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 0.1 % Triton X-100, (pH 7.5 @ 25°C)
Specification Version: PS-M0545S/L v1.0
Effective Date: 22 May 2018

Assay Name/Specification (minimum release criteria)	Lot #0031708
Endonuclease Activity (Circular Single Stranded DNA) - A 50 µl reaction in NEBuffer 4 containing 1 µg of M13mp18 Single-stranded DNA and a minimum of 30 units of Exonuclease VIII, truncated incubated for 4 hours at 37°C results in <10% conversion to linear DNA as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicked Circular DNA) - A 50 µl reaction in NEBuffer 4 containing 1 µg of PhiX174 RF II DNA and a minimum of 30 units of Exonuclease VIII, truncated incubated for 4 hours at 37°C results in <10% conversion to linear DNA as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking) - A 50 µl reaction in NEBuffer 4 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of Exonuclease VIII, truncated incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) - Exonuclease VIII, truncated is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 10 units of Exonuclease VIII, truncated is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass



Authorized by
Derek Robinson
22 May 2018



Inspected by
John Greci
24 Aug 2017

