

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: EnGen® Spy Cas9 Nickase

Catalog Number: M0650T
Concentration: 20 µM
Unit Definition: N/A

Packaging Lot Number: 10102988
Expiration Date: 03/2023
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl , 300 mM NaCl , 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0650T v1.0

EnGen® Spy Cas9 Nickase Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0650TVIAL	EnGen® Spy Cas9 Nickase	10102989	Pass	
B6003SVIAL	NEBuffer™ r3.1	10102967	Pass	

Assay Name/Specification	Lot # 10102988
Functional Testing (EnGen® Spy Cas9 Nickase, Targeted Nicking) A 20 µl reaction in 1X NEBuffer 3.1 containing 20 nM 100 bp FAM and ROX labeled double stranded target DNA, 100 nM sgRNA and 100 nM EnGen® Spy Cas9 Nickase incubated for 1 hour at 37°C results in ≥90% nicking of the substrate DNA as determined by capillary electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 1 pmol of EnGen® Spy Cas9 Nickase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 3.1 containing 1 μg of supercoiled PhiX174 RF I DNA and a minimum of 1 pmol of EnGen® Spy Cas9 Nickase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 pmol of EnGen® Spy Cas9 Nickase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by	Pass



M0650T / Lot: 10102988

Page 1 of 2

Assay Name/Specification	Lot # 10102988
gel electrophoresis using fluorescent detection.	
Protein Purity Assay (SDS-PAGE) EnGen® Spy Cas9 Nickase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 1 pmol of EnGen® Spy Cas9 Nickase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Bhairavi Jani Production Scientist

06 May 2021



Packaging Quality Control Inspector

06 May 2021



M0650T / Lot: 10102988

Page 2 of 2