

New England Biolabs Certificate of Analysis

Product Name: EnGen® Spy dCas9 (SNAP-tag)
Catalog Number: M0652S
Concentration: 1,000 nM
Unit Definition: N/A
Packaging Lot Number: 10081025
Expiration Date: 08/2022
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl , 300 mM NaCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version: PS-M0652S v1.0

EnGen® Spy dCas9 (SNAP-tag) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0652SVIAL	EnGen® Spy dCas9 (SNAP-tag®)	10081026	Pass
B7203SVIAL	NEBuffer™ 3.1	10077593	Pass

Assay Name/Specification	Lot # 10081025
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 1 pmol of EnGen® Spy dCas9 (SNAP-tag®) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 1 pmol of EnGen® Spy dCas9 (SNAP-tag®) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (EnGen® Spy dCas9 (SNAP-tag®), Gel Shift Assay) A 20 µl reaction in 1X NEBuffer 3.1 containing 20 nM 100 bp FAM labeled double stranded target DNA, 20 nM TAMRA-labeled off target DNA, 100 nM sgRNA and 100 nM EnGen® Spy dCas9 (SNAP-tag®) incubated for 15 minutes at 37°C results in ≥90% binding of the substrate DNA as determined by electrophoretic mobility shift assay.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 1 pmol of EnGen® Spy dCas9 (SNAP-tag®) incubated for 16 hours at 37°C results in a DNA	Pass

Assay Name/Specification	Lot # 10081025
<p>pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	
<p>Protein Purity Assay (SDS-PAGE) EnGen[®] Spy dCas9 (SNAP-tag[®]) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 pmol of EnGen[®] Spy dCas9 (SNAP-tag[®]) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Bhairavi Jani
Production Scientist
07 Aug 2020



Josh Hersey
Packaging Quality Control Inspector
07 Aug 2020