

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: EnGen Lba Cas12a (Cpf1)

Catalog Number:M0653TConcentration:100 μMPackaging Lot Number:10119060Expiration Date:03/2024Storage Temperature:-20°C

Storage Conditions: 500 mM NaCl, 20 mM Sodium Acetate, 0.1 mM EDTA, 0.1 mM TCEP-HCl, 50%

Glycerol, (pH 6.0 @ 25°C)

Specification Version: PS-M0653T v2.0

EnGen Lba Cas12a (Cpf1) Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0653TVIAL	EnGen® Lba Cas12a (Cpf1)	10130322	Pass	
B6002SVIAL	NEBuffer™ r2.1	10103928	Pass	
B0653AVIAL	EnGen® Lba Cas12a Diluent	10122103	Pass	

Assay Name/Specification	Lot # 10119060
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2.1 containing 1 µg of Lambda DNA and a minimum of 1 pmol of EnGen® Lba Cas12a (Cpf1) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 2.1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 1 pmol of EnGen® Lba Cas12a (Cpf1) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (Targeted Digestion) A 20 µl reaction in 1X NEBuffer 2.1 containing 20 nM of 100 bp FAM and ROX-labeled double-stranded target DNA, 100 nM crRNA, and 100 nM EnGen® Lba Cas12a (Cpf1) incubated for 15 minutes at 37°C results in ≥90% targeted digestion of the substrate DNA as determined by capillary electrophoresis.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 2.1 containing 1 μg of supercoiled PhiX174 RF I DNA and a minimum of 1 pmol of EnGen® Lba Cas12a (Cpf1) incubated for 4 hours at 37°C	Pass



M0653T / Lot: 10119060

Page 1 of 2

Assay Name/Specification	Lot # 10119060
results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of f-300 RNA transcript and a minimum of 1 pmol of EnGen® Lba Cas12a (Cpf1) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Bhairavi Jani Production Scientist 25 Mar 2022 Michael Tonello

Packaging Quality Control Inspector

25 Mar 2022



M0653T / Lot: 10119060 Page 2 of 2