

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: 5-hydroxythymidine DNA Kinase

Catalog Number: M0659S Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to protect 1 µg

of Bacillus subtilis bacteriophage SP8 genomic DNA in ##30 minutes

at 37°C in a total reaction volume of 20 µl against cleavage by Ncol-HF restriction endonuclease.

Packaging Lot Number: 10156687 Expiration Date: 08/2024

Storage Temperature:

Storage Conditions: 10 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol,

(pH 7.4 @ 25°C)

-20°C

Specification Version: PS-M0659S v1.0

5-hydroxythymidine DNA Kinase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0659SVIAL	5-hydroxymethyluridine DNA Kinase	10156686	Pass	
B0202SVIAL	T4 DNA Ligase Reaction Buffer	10153862	Pass	

Assay Name/Specification	Lot # 10156687
Endonuclease Activity (Nicking)	Pass
A 50 μl reaction in T4 DNA Ligase Reaction Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 20 units of 5-hydroxymethyluridine DNA Kinase incubated	
for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in T4 DNA Ligase Reaction Buffer containing 1 µg of Lambda DNA and a minimum of 20 units of 5-hydroxymethyluridine DNA Kinase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in T4 DNA Ligase Reaction Buffer containing 1 µg of a mixture of	
single and double-stranded [³H] E. coli DNA and a minimum of 20 units of	
5-hydroxymethyluridine DNA Kinase incubated for 4 hours at 37°C releases <0.1% of	
the total radioactivity.	



M0659S / Lot: 10156687

Page 1 of 2

Assay Name/Specification	Lot # 10156687
	_
Protein Purity Assay (SDS-PAGE)	Pass
5-hydroxymethyluridine DNA Kinase is ≥ 95% pure as determined by SDS-PAGE analysis	
using Coomassie Blue detection.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Jenna Ware Production Scientist

31 Aug 2022

Erin Varney

Packaging Quality Control Inspector

31 Aug 2022

