

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Tte-UvrD Helicase

Catalog Number: M1202S
Concentration: 20 µg/ml
Packaging Lot Number: 10221166
Expiration Date: 11/2025
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.1 %

Triton®X-100 , 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M1202S v1.0

Tte-UvrD Helicase Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
P0756SVIAL	Adenosine 5'-Triphosphate (ATP)	10211010	Pass	
M1202SVIAL	Tte-UvrD Helicase	10221167	Pass	
B0537SVIAL	Isothermal Amplification Buffer	10201499	Pass	

Assay Name/Specification	Lot # 10221166
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart® Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 2 μg of Tte-UvrD Helicase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart® Buffer containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 2 μg of Tte-UvrD Helicase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (ATP-dependent LAMP Inhibition) Tte-UvrD Helicase is tested for ATP-dependent inhibition of a LAMP reaction and suppression of amplification in a no template control reaction.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 2 µg of Tte-UvrD Helicase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass



M1202S / Lot: 10221166

Page 1 of 2

Assay Name/Specification	Lot # 10221166 Pass
Protein Purity Assay (SDS-PAGE)  Tte-UvrD Helicase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Tte-UvrD Helicase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Alicia Bielik Production Scientist 22 Nov 2023 Josh Hersey
Packaging Quality (

Packaging Quality Control Inspector 29 Dec 2023



M1202S / Lot: 10221166

Page 2 of 2