

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Hi-T4 DNA Ligase

Catalog Number: M2622L
Concentration: 400,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to give 50%

ligation of 6 µg of Lambda-HindIII DNA in 30 minutes at 25°C in a

total reaction volume of 20 µl.

Packaging Lot Number: 10058735 Expiration Date: 11/2021 Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol,

(pH 7.4 @ 25°C)

Specification Version: PS-M2622S/L v1.0

Hi-T4 DNA Ligase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M2622LVIAL	Hi-T4 DNA Ligase	10058610	Pass	
B0535AVIAL	StickTogether™ DNA Ligase Buffer	10053903	Pass	
B0202SVIAL	T4 DNA Ligase Reaction Buffer	10041768	Pass	

Assay Name/Specification	Lot # 10058735
Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 2000 units of Hi-T4™ DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Hi-T4™ DNA Ligase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 400 units of Hi-T4™ DNA Ligase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli	Pass



M2622L / Lot: 10058735 Page 1 of 3

Assay Name/Specification	Lot # 10058735
genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	
Protein Purity Assay (SDS-PAGE) Hi-T4™ DNA Ligase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Protein Concentration (A280) The concentration of Hi-T4™ DNA Ligase is 0.4 mg/ml +/- 10% as determined by UV absorption at 280 nm. Protein concentration is determined by the Pace method using the extinction coefficient of 57,675 and molecular weight of 56,806 daltons for Hi-T4™ DNA Ligase (Pace, C.N. et al. (1995) Protein Sci., 4, 2411-2423).	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 1 containing 1 µg of CIP-treated Lambda-HindIII DNA and a minimum of 400 units of Hi-T4™ DNA Ligase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 400 units of Hi-T4™ DNA Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Double Stranded DNase Activity (Labeled Oligo) A 50 μl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a blunt end and a minimum of 2000 units of Hi-T4™ DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
DNase Activity (Labeled Oligo, 5' extension) A 50 μl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent abeled double-stranded oligonucleotide containing a 5' extension and a minimum of 2000 units of Hi-T4™ DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
DNase Activity (Labeled Oligo, 3' extension) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 3' extension and a minimum of 2000 units of Hi-T4™ DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.



M2622L / Lot: 10058735 Page 2 of 3



May K brenzen

Mary Lorenzen
Production Scientist
09 Sep 2019

Michael Tonello

Packaging Quality Control Inspector

11 Nov 2019