

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Hi-T4™ DNA Ligase

Catalog Number: M2622L
Concentration: 400,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to give 50%

ligation of 6 µg of Lambda-HindIII DNA in 30 minutes at 25°C in a

total reaction volume of 20 μl.

Packaging Lot Number: 10097783
Expiration Date: 12/2022
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol,

(pH 7.4 @ 25°C)

Specification Version: PS-M2622S/L v1.0

| Hi-T4™ DNA Ligase Component List | | | | |
|----------------------------------|----------------------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| M2622LVIAL | Hi-T4™ DNA Ligase | 10092877 | Pass | |
| B0535AVIAL | StickTogether™ DNA Ligase Buffer | 10077408 | Pass | |
| B0202SVIAL | T4 DNA Ligase Reaction Buffer | 10088347 | Pass | |

| Assay Name/Specification | Lot # 10097783 |
|---|----------------|
| DNase Activity (Labeled Oligo, 5' extension) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 5' extension and a minimum of 2000 units of Hi-T4™ DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 μl reaction in NEBuffer 1 containing 1 μg of CIP-treated Lambda-HindIII DNA and a minimum of 400 units of Hi-T4 [™] DNA Ligase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Protein Concentration (A280) The concentration of Hi-T4™ DNA Ligase is 0.4 mg/ml +/- 10% as determined by UV absorption at 280 nm. Protein concentration is determined by the Pace method using the extinction coefficient of 57,675 and molecular weight of 56,806 daltons for | Pass |



M2622L / Lot: 10097783

Page 1 of 3

| Assay Name/Specification | Lot # 10097783 |
|--|----------------|
| Hi-T4™ DNA Ligase (Pace, C.N. et al. (1995) Protein Sci., 4, 2411-2423). | |
| Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 400 units of Hi-T4™ DNA Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Protein Purity Assay (SDS-PAGE) Hi-T4™ DNA Ligase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection. | Pass |
| Double Stranded DNase Activity (Labeled Oligo) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a blunt end and a minimum of 2000 units of Hi-T4™ DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis. | Pass |
| DNase Activity (Labeled Oligo, 3' extension) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 3' extension and a minimum of 2000 units of Hi-T4™ DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis. | Pass |
| Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 2000 units of Hi-T4™ DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis. | Pass |
| RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Hi-T4™ DNA Ligase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection. | Pass |
| qPCR DNA Contamination (E. coli Genomic) A minimum of 400 units of Hi-T4™ DNA Ligase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome. | Pass |

This product has been tested and shown to be in compliance with all specifications.



M2622L / Lot: 10097783

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ana Egana **Production Scientist**

NEW ENGLAND

09 Feb 2021

Josh Hersey Packaging Quality Control Inspector

09 Feb 2021



M2622L / Lot: 10097783

Page 3 of 3