

New England Biolabs Certificate of Analysis

Product Name: Deoxynucleotide (dNTP) Solution Set
Catalog Number: N0446S
Concentration: 100 mM
Unit Definition: N/A
Packaging Lot Number: 10256838
Expiration Date: 01/2026
Storage Temperature: -20°C
Storage Conditions: Supplied in Ultrapure water as a sodium salt (pH 7.5)
Specification Version: PS-N0446S/V v3.0


Deoxynucleotide (dNTP) Solution Set Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
N0443SVIAL	dTTP	10226741	Pass
N0442SVIAL	dGTP	10226738	Pass
N0441SVIAL	dCTP	10226736	Pass
N0440SVIAL	dATP Solution	10226710	Pass

Assay Name/Specification	Lot # 10256838
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 2 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 1 µl of dATP, dCTP, dGTP, and dTTP incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 4 µl of dATP, dCTP, dGTP, and dTTP incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
PCR Amplification (0.5 kb Lambda, dNTPs) A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM dATP, dCTP, dGTP, and dTTP and 0.2 µM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 0.5 kb product.	Pass
PCR Amplification (2.0 kb Lambda, dNTPs) A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM dATP, dCTP,	Pass

Assay Name/Specification	Lot # 10256838
<p>dGTP, and dTTP and 0.2 µM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 2.0 kb product.</p>	
<p>PCR Amplification (5.0 kb Lambda, dNTPs) A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM dATP, dCTP, dGTP, and dTTP and 0.2 µM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.</p>	Pass
<p>Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 16 µl of dATP, dCTP, dGTP, and dTTP incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	Pass
<p>Physical Purity (HPLC) dATP, dCTP, dGTP, and dTTP is ≥ 99% pure as determined by HPLC analysis.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl dATP, dCTP, dGTP, and dTTP is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 1 µl of dATP, dCTP, dGTP, and dTTP is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Lea Antonopoulos
Production Scientist
04 Apr 2024



Michael Tonello
Packaging Quality Control Inspector
29 Aug 2024