240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350

www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

TriDye<sup>TM</sup> 2-Log DNA Ladder (0.1 - 10.0 kb) Product Name:

N3270S Catalog #: Concentration:  $100 \,\mu\mathrm{g/ml}$ Unit Definition: N/A*Lot* #: 0101505 05/2015 Assay Date: Expiration Date: 5/2017 4°C Storage Temp:

0.006 % Xylene cyanol, 10 mM Tris-HCl (pH 7.9), 10 mM EDTA, 10 % Glycerol, 0.006 % Bromophenol Blue, 0.06 % Storage Conditions:

Orange G

PS-N3270S v1.0 Specification Version: Effective Date: 25 Jun 2014

Assay Name/Specification (minimum release criteria)	Lot #0101505
<b>DNA Concentration (A260)</b> - The concentration of TriDye <sup>TM</sup> 2-Log DNA Ladder (0.1 - 10.0 kb) is between 100 and 110 μg/ml as determined by UV absorption at 260 nm.	Pass
<b>Electrophoretic Pattern (Marker)</b> - The banding pattern of TriDye <sup>TM</sup> 2-Log DNA Ladder (0.1 - 10.0 kb) on a 1.2% agarose gel shows discrete, clearly identifiable bands at each band of the marker, when stained with Ethidium Bromide at a concentration of 0.5 μg/ml.	Pass
Non-Specific DNase Activity (DNA, 16 hour) - A 50 µl reaction in 1X NEBuffer 2 containing 5 µg of TriDye TM 2-Log DNA Ladder (0.1 - 10.0 kb) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

Authorized by Derek Robinson 25 Jun 2014







Vanessa Mathieu-Sheltry

05 May 2015