

## New England Biolabs Certificate of Analysis

*Product Name:* PNGase F, Recombinant  
*Catalog #:* P0708S/L  
*Concentration:* 500,000 units/ml  
*Unit Definition:* One unit is defined as the amount of enzyme required to remove > 95% of the carbohydrate from 10 µg of denatured RNase B in 1 hour at 37°C in a total reaction volume of 10 µl. (65 NEB units = 1 IUB milliunit).  
*Lot #:* 0021611  
*Assay Date:* 11/2016  
*Expiration Date:* 11/2018  
*Storage Temp:* -20°C  
*Storage Conditions:* 50 mM NaCl , 20 mM Tris-HCl , 5 mM EDTA , 50 % Glycerol, (pH 7.5 @ 25°C)  
*Specification Version:* PS-P0708S/L v1.0  
*Effective Date:* 17 Feb 2016

Assay Name/Specification (minimum release criteria)	Lot #0021611
<b>Glycosidase Activity (Endo F1, F2, H)</b> - A 10 µl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>
<b>Glycosidase Activity (Endo F2, F3)</b> - A 10 µl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>
<b>Glycosidase Activity (β-Mannosidase)</b> - A 10 µl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>
<b>Glycosidase Activity (β-Xylosidase)</b> - A 10 µl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>
<b>Glycosidase Activity (β1-3 Galactosidase)</b> - A 10 µl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>

## New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0021611
<b>Glycosidase Activity (<math>\beta</math>1-4 Galactosidase)</b> - A 10 $\mu$ l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled $\beta$ -Galactosidase substrate (Gal $\beta$ 1-4GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc -AMC) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>
<b>Glycosidase Activity (<math>\beta</math>-N-Acetylgalactosaminidase)</b> - A 10 $\mu$ l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled $\beta$ -N-Acetylgalactosaminidase substrate (GalNAc $\beta$ 1-4Gal $\beta$ 1-4Glc-AMC) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>
<b>Glycosidase Activity (<math>\beta</math>-N-Acetylglucosaminidase)</b> - A 10 $\mu$ l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled $\beta$ -N-Acetylglucosaminidase substrate (GlcNAc $\beta$ 1-4GlcNAc $\beta$ 1-4GlcNAc-AMC) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>
<b>Glycosidase Activity (<math>\alpha</math>-Glucosidase)</b> - A 10 $\mu$ l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled $\alpha$ -Glucosidase substrate (Glc $\alpha$ 1-6Glc $\alpha$ 1-4Glc-AMC) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>
<b>Glycosidase Activity (<math>\alpha</math>-Neuraminidase)</b> - A 10 $\mu$ l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled $\alpha$ -Neuraminidase substrate (Neu5Ac $\alpha$ 2-3Gal $\beta$ 1-3GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>
<b>Glycosidase Activity (<math>\alpha</math>1-2 Fucosidase)</b> - A 10 $\mu$ l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled $\alpha$ -Fucosidase substrate (Fuc $\alpha$ 1-2Gal $\beta$ 1-4Glc-AMC) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>
<b>Glycosidase Activity (<math>\alpha</math>1-3 Fucosidase)</b> - A 10 $\mu$ l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled $\alpha$ -Fucosidase substrate (Fuc $\alpha$ 1-3Gal $\beta$ 1-4GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-AMC) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>
<b>Glycosidase Activity (<math>\alpha</math>1-3 Galactosidase)</b> - A 10 $\mu$ l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled $\alpha$ -Galactosidase substrate (Gal $\alpha$ 1-3Gal $\beta$ 1-4GlcNAc-AMC) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	<b>Pass</b>

## New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0021611
<p><b>Glycosidase Activity (<math>\alpha</math>1-3 Mannosidase)</b> - A 10 <math>\mu</math>l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Mannosidase substrate (Man<math>\alpha</math>1-3Man<math>\beta</math>1-4GlcNAc-AMC) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>1-6 Galactosidase)</b> - A 10 <math>\mu</math>l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Galactosidase substrate (Gal<math>\alpha</math>1-6Gal<math>\alpha</math>1-6Glc<math>\alpha</math>1-2Fru-AMC) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>1-6 Mannosidase)</b> - A 10 <math>\mu</math>l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Mannosidase substrate (Man<math>\alpha</math>1-6Man<math>\alpha</math>1-6(Man<math>\alpha</math>1-3)Man-AMC) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>-N-Acetylgalactosaminidase)</b> - A 10 <math>\mu</math>l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled <math>\alpha</math>-N-Acetylgalactosaminidase substrate (GalNAc<math>\alpha</math>1-3(Fuc<math>\alpha</math>1-2)Gal<math>\beta</math>1-4Glc-AMC) and 5,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Protease Activity (SDS-PAGE)</b> - A 20 <math>\mu</math>l reaction in 1X Glyco Buffer 2 containing 24 <math>\mu</math>g of a standard mixture of proteins and a minimum of 10,000 units of PNGase F, Recombinant incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>Protein Purity Assay (SDS-PAGE)</b> - PNGase F, Recombinant is <math>\geq</math> 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>



Authorized by  
Derek Robinson  
17 Feb 2016



Inspected by  
Alicia Bielik  
07 Nov 2016

