

## New England Biolabs Certificate of Analysis

**Product Name:** Rapid™ PNGase F (non-reducing format)  
**Catalog Number:** P0711S  
**Unit Definition:** N/A  
**Packaging Lot Number:** 10124414  
**Expiration Date:** 10/2022  
**Storage Temperature:** 4°C  
**Storage Conditions:** 50 mM NaCl , 20 mM Tris-HCl , 5 mM EDTA, (pH 7.5 @ 25°C)  
**Specification Version:** PS-P0711S v2.0

Rapid™ PNGase F (non-reducing format) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P0711SVIAL	Rapid™ PNGase F (non-reducing format)	10124413	Pass
B0717SVIAL	5X Rapid PNGase F Buffer (non-reducing format)	10124415	Pass

Assay Name/Specification	Lot # 10124414
<b>Glycosidase Activity (β-N-Acetylgalactosaminidase)</b> A 10 µl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 1 µl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
<b>Glycosidase Activity (β-N-Acetylglucosaminidase)</b> A 10 µl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 1 µl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
<b>Glycosidase Activity (β-Xylosidase)</b> A 10 µl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 1 µl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
<b>Glycosidase Activity (α1-3 Fucosidase)</b> A 10 µl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of	Pass

Assay Name/Specification	Lot # 10124414
<p>fluorescently-labeled <math>\alpha</math>-Fucosidase substrate (Fuc<math>\alpha</math>1-3Gal<math>\beta</math>1-4GlcNAc<math>\beta</math>1-3Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p><b>Glycosidase Activity (<math>\alpha</math>1-3 Galactosidase)</b> A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-Galactosidase substrate (Gal<math>\alpha</math>1-3Gal<math>\beta</math>1-4GlcNAc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>1-2 Fucosidase)</b> A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-Fucosidase substrate (Fuc<math>\alpha</math>1-2Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Protease Activity (SDS-PAGE)</b> A 20 <math>\mu</math>l reaction in 1X Rapid PNGase F Buffer (non-reducing format) containing 24 <math>\mu</math>g of a standard mixture of proteins and a minimum of 5 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>Protein Purity Assay (SDS-PAGE)</b> Rapid PNGase F (non-reducing format) is <math>\geq</math> 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (Endo F2, F3)</b> A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (Endo F1, F2, H)</b> A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>-Neuraminidase)</b> A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-Neuraminidase substrate</p>	<b>Pass</b>

Assay Name/Specification	Lot # 10124414
<p>(Neu5Ac<math>\alpha</math>2-3Gal<math>\beta</math>1-3GlcNAc<math>\beta</math>1-3Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p><b>Glycosidase Activity (<math>\alpha</math>-N-Acetylgalactosaminidase)</b> A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-N-Acetylgalactosaminidase substrate (GalNAc<math>\alpha</math>1-3(Fuc<math>\alpha</math>1-2)Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>-Glucosidase)</b> A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-Glucosidase substrate (Glc<math>\alpha</math>1-6Glc<math>\alpha</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>1-6 Mannosidase)</b> A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-Mannosidase substrate (Man<math>\alpha</math>1-6Man<math>\alpha</math>1-6(Man<math>\alpha</math>1-3)Man-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>1-3 Mannosidase)</b> A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-Mannosidase substrate (Man<math>\alpha</math>1-3Man<math>\beta</math>1-4GlcNAc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>1-6 Galactosidase)</b> A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-Galactosidase substrate (Gal<math>\alpha</math>1-6Gal<math>\alpha</math>1-6Glc<math>\alpha</math>1-2Fru-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\beta</math>-Mannosidase)</b> A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\beta</math>-Mannosidase substrate (Man<math>\beta</math>1-4Man<math>\beta</math>1-4Man-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\beta</math>1-3 Galactosidase)</b> A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of</p>	<b>Pass</b>

Assay Name/Specification	Lot # 10124414
<p>fluorescently-labeled <math>\beta</math>-Galactosidase substrate (Gal<math>\beta</math>1-3GlcNAc<math>\beta</math>1-4Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> <p><b>Glycosidase Activity (<math>\beta</math>1-4 Galactosidase)</b> A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\beta</math>-Galactosidase substrate (Gal<math>\beta</math>1-4GlcNAc<math>\beta</math>1-3Gal<math>\beta</math>1-4Glc -AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Alicia Bielik  
Production Scientist  
14 Jan 2022



Michael Tonello  
Packaging Quality Control Inspector  
14 Jan 2022