

New England Biolabs Certificate of Analysis

Product Name: α 1-2,3 Mannosidase
Catalog Number: P0729S
Concentration: 32,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to cleave > 95% of the non-reducing terminal α -D-mannose from 1 nmol Man α 1-3Man β 1-4GlcNAc-7-amino-4-methyl-coumarin (AMC), in 1 hour at 37°C in a total reaction volume of 10 μ l.
Packaging Lot Number: 10092475
Expiration Date: 12/2021
Storage Temperature: 4°C
Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, (pH 7.5 @ 25°C)
Specification Version: PS-P0729S/L v1.0

α1-2,3 Mannosidase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P0729SVIAL	α 1-2,3 Mannosidase	10092474	Pass
B9001SVIAL	Purified BSA	10098897	Pass
B1727SVIAL	10X GlycoBuffer 1	10092862	Pass

Assay Name/Specification	Lot # 10092475
Glycosidase Activity (β-Xylosidase) A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β -Xylosidase substrate (Xyl β 1-4Xyl β 1-4Xyl β 1-4Xyl-AMC) and 32 units of α 1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-3 Galactosidase) A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β -Galactosidase substrate (Gal β 1-3GlcNAc β 1-4Gal β 1-4Glc-AMC) and 32 units of α 1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-4 Galactosidase) A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β -Galactosidase substrate (Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc -AMC) and 32 units of α 1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass

Assay Name/Specification	Lot # 10092475
<p>Protease Activity (SDS-PAGE) A 20 µl reaction in 1X Glyco Buffer 1 containing 24 µg of a standard mixture of proteins and a minimum of 400 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) α1-2,3 Mannosidase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>Glycosidase Activity (Endo F1, F2, H) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (Endo F2, F3) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (PNGase F) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled PNGase F substrate (Fluoresceinated fetuin triantennary) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Glucosidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Glucosidase substrate (Glcα1-6Glcα1-4Glc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-N-Acetylgalactosaminidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-N-Acetylgalactosaminidase substrate (GalNAcα1-3(Fucα1-2)Galβ1-4Glc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Neuraminidase)</p>	Pass

Assay Name/Specification	Lot # 10092475
<p>A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Neuraminidase substrate (Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p>Glycosidase Activity (α1-2 Fucosidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fuca1-2Galβ1-4Glc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Fucosidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fuca1-3Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Galactosidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-3Galβ1-4GlcNAc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-6 Galactosidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-6Galα1-6Glcα1-2Fru-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-Mannosidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-N-Acetylgalactosaminidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-N-Acetylglucosaminidase) A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled</p>	Pass

Assay Name/Specification	Lot # 10092475
<p>β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Alicia Bielik
Production Scientist
23 Feb 2021



Michael Tonello
Packaging Quality Control Inspector
23 Feb 2021