

## New England Biolabs Certificate of Analysis


**Product Name:** *Bacteroides Heparinase I*  
**Catalog Number:** *P0735L*  
**Concentration:** *12,000 U/ml*  
**Unit Definition:** *One unit is defined as the amount of enzyme that will liberate 1.0  $\mu$ mol unsaturated oligosaccharides from porcine mucosal heparin per minute at 30°C and pH 7.0 in a total reaction volume of 100  $\mu$ l.*  
**Lot Number:** *10051176*  
**Expiration Date:** *07/2020*  
**Storage Temperature:** *-80°C*  
**Storage Conditions:** *100 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, 5 mM CaCl<sub>2</sub>, (pH 7.5 @ 25°C)*  
**Specification Version:** *PS-P0735S/L v1.0*

Bacteroides Heparinase I Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P0735LVIAL	Bacteroides Heparinase I	10050305	Pass
B0735SVIAL	Bacteroides Heparinase Reaction Buffer (10X)	10034984	Pass

Assay Name/Specification	Lot # 10051176
<p><b>Protease Activity (SDS-PAGE)</b>            A 20 <math>\mu</math>l reaction in 1X Heparinase Reaction Buffer containing 24 <math>\mu</math>g of a standard mixture of proteins and a minimum of 120 units of Bacteroides Heparinase I incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>Protein Purity Assay (SDS-PAGE)</b>            Bacteroides Heparinase I is <math>\geq</math> 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>Sulfatase Activity (2-O)</b>            A 10 <math>\mu</math>l reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled 2-O-Sulfatase substrate (<math>\Delta</math>UA2S-(1-4)-GlcNS6S-AMC) and 24 units of Bacteroides Heparinase I incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Sulfatase and Uronidase Activity (N,6-O)</b>            A 10 <math>\mu</math>l reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled N,6-O-Sulfatase substrate (<math>\Delta</math>UA-(1-4)-GlcNS6S-AMC) and 24 units</p>	<b>Pass</b>

Assay Name/Specification	Lot # 10051176
<p>of Bacteroides Heparinase I incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p><b>Glycosidase Activity (β1-3 Galactosidase)</b> A 10 µl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 24 units of Bacteroides Heparinase I incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (β1-4 Galactosidase)</b> A 10 µl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc -AMC) and 24 units of Bacteroides Heparinase I incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (β-N-Acetylgalactosaminidase)</b> A 10 µl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 24 units of Bacteroides Heparinase I incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (β-N-Acetylglucosaminidase)</b> A 10 µl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 24 units of Bacteroides Heparinase I incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.



Brad Landgraf  
Production Scientist  
24 Jan 2019



Michael Tonello  
Packaging Quality Control Inspector  
05 Aug 2019