

New England Biolabs Certificate of Analysis


Product Name: *Bacteroides Heparinase II*
Catalog Number: P0736S
Concentration: 4,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme that will liberate 1.0 μ mol unsaturated oligosaccharides from porcine mucosal heparin per minute at 30°C and pH 7.0 in a total reaction volume of 100 μ l.
Lot Number: 10030741
Expiration Date: 12/2019
Storage Temperature: -80°C
Storage Conditions: 100 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, 5 mM CaCl₂, (pH 7.5 @ 25°C)
Specification Version: PS-P0736S/L v1.0

Bacteroides Heparinase II Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P0736SVIAL	Bacteroides Heparinase II	10030742	Pass
B0735SVIAL	Bacteroides Heparinase Reaction Buffer (10X)	0071803	Pass


Assay Name/Specification	Lot # 10030741
<p>Glycosidase Activity (β1-3 Galactosidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 8 units of Bacteroides Heparinase II incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β1-4 Galactosidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 8 units of Bacteroides Heparinase II incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-N-Acetylgalactosaminidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 8 units of Bacteroides Heparinase II incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass

Assay Name/Specification	Lot # 10030741
<p>Glycosidase Activity (β-N-Acetylglucosaminidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 8 units of Bacteroides Heparinase II incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Protease Activity (SDS-PAGE) A 20 μl reaction in 1X Heparinase Reaction Buffer containing 24 μg of a standard mixture of proteins and a minimum of 20 units of Bacteroides Heparinase II incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) Bacteroides Heparinase II is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>Sulfatase Activity (2-O) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled 2-O-Sulfatase substrate (ΔUA2S-(1-4)-GlcNS6S-AMC) and 8 units of Bacteroides Heparinase II incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Sulfatase and Uronidase Activity (N,6-O) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled N,6-O-Sulfatase substrate (ΔUA-(1-4)-GlcNS6S-AMC) and 8 units of Bacteroides Heparinase II incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Brad Landgraf
Production Scientist
05 Dec 2018



Josh Hersey
Packaging Quality Control Inspector
07 Dec 2018