

## New England Biolabs Certificate of Analysis

**Product Name:** Adenosine-5 Triphosphate (ATP)  
**Catalog Number:** P0756L  
**Concentration:** 10 mM  
**Unit Definition:** N/A  
**Packaging Lot Number:** 10118110  
**Expiration Date:** 07/2023  
**Storage Temperature:** -20°C  
**Storage Conditions:** Milli-Q® Water as a sodium salt, (pH 7.0 @ 25°C)  
**Specification Version:** PS-P0756S/L v2.0

Adenosine-5 Triphosphate (ATP) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P0756SVIAL	Adenosine 5'-Triphosphate (ATP)	10109057	Pass

Assay Name/Specification	Lot # 10118110
<b>RNase Activity (Extended Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 mM of ATP is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
<b>Protease Activity (SDS-PAGE)</b> A 20 µl reaction in 1X CutSmart® Buffer containing 24 µg of a standard mixture of proteins and a minimum of 1 mM of ATP incubated for 16 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.	Pass
<b>Phosphatase Activity (pNPP)</b> A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl <sub>2</sub> containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 1 mM of ATP incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 1 mM of ATP incubated for 4 hours at 30°C releases <0.1% of the total radioactivity.	Pass

Assay Name/Specification	Lot # 10118110
<p><b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 1 mM of ATP incubated for 4 hours at 30°C results in &lt;10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 10 µl of ATP incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Alicia Bielik  
Production Scientist  
03 Aug 2021



Michael Tonello  
Packaging Quality Control Inspector  
03 Aug 2021