

## New England Biolabs Certificate of Analysis

**Product Name:** *Stul*

**Catalog Number:** *R0187L*

**Concentration:** *10,000 U/ml*

**Unit Definition:** *One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.*

**Packaging Lot Number:** *10156872*

**Expiration Date:** *07/2024*

**Storage Temperature:** *-20°C*

**Storage Conditions:** *10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)*

**Specification Version:** *PS-R0187S/L v2.0*

Stul Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0187LVIAL	Stul	10156871	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10153339	Pass
B6004SVIAL	rCutSmart™ Buffer	10154051	Pass

Assay Name/Specification	Lot # 10156872
<p><b>Blue-White Screening (Terminal Integrity)</b> A sample of Litmus 28i vector linearized with a 10-fold excess of Stul, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in &lt;1% white colonies.</p>	Pass
<p><b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of Lambda DNA with Stul, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with Stul.</p>	Pass
<p><b>Protein Purity Assay (SDS-PAGE)</b> Stul is &gt;95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.</p>	Pass
<p><b>Functional Testing (15 minute Digest)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 µl of Stul incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.</p>	Pass

Assay Name/Specification	Lot # 10156872
<p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of Stul incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 10 units of Stul is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	<b>Pass</b>
<p><b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled pBR322 DNA and a minimum of 10 units of Stul incubated for 4 hours at 37°C results in &lt;20% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 300 units of Stul incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Penghua Zhang  
Production Scientist  
05 Aug 2022



Michael Tonello  
Packaging Quality Control Inspector  
05 Aug 2022