

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	AfIII
Catalog Number:	R0520L
Concentration:	20,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of PhiX174 RF I DNA in 1 hour at 37°C in a total reaction volume of 50 μl.
Lot Number:	10013033
Expiration Date:	06/2020
Storage Temperature:	-20°C
Storage Conditions:	50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA
Specification Version:	PS-R0520S/L v1.0

AfIII Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0520LVIAL	AfIII	10013034	Pass	
B7204SVIAL	CutSmart® Buffer	10010634	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10011266	Pass	

Assay Name/Specification	Lot # 10013033
Ligation and Recutting (Terminal Integrity) After a 5-fold over-digestion of PhiX174 DNA with AfIII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AfIII.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of PhiX174 DNA and a minimum of 200 Units of AfIII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Blue-White Screening (Terminal Integrity) A sample of Litmus28 vector linearized with a 10-fold excess of AfIII, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled pBR322 DNA and a	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10013033
minimum of 100 Units of AfIII incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	
double-stranded [³ H] E. coli DNA and a minimum of 100 units of AfIII incubated for	
4 hours at 37°C releases <0.1% of the total radioactivity.	

This product has been tested and shown to be in compliance with all specifications.

Yor

Tony Spear-Alfonso Production Scientist 06 Jun 2018

Michae Mr.

Michael Tonello Packaging Quality Control Inspector 24 Jul 2018

