

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	AfIII
Catalog Number:	R0520S
Concentration:	20,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μ g of PhiX174 RF I DNA in 1 hour at 37°C in a total reaction volume of 50 μ l.
Lot Number:	10013035
Expiration Date:	06/2020
Storage Temperature:	-20°C
Storage Conditions:	50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA
Specification Version:	PS-R0520S/L v1.0

AfIII Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0520SVIAL	AfIII	10013036	Pass	
B7204SVIAL	CutSmart® Buffer	3091805	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	0251805	Pass	

Assay Name/Specification	Lot # 10013035
Blue-White Screening (Terminal Integrity) A sample of Litmus28 vector linearized with a 10-fold excess of AfIII, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled pBR322 DNA and a minimum of 100 Units of AfIII incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of AfIII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 5-fold over-digestion of PhiX174 DNA with AfIII, >95% of the DNA fragments	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10013035
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AfIII.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of PhiX174 DNA and a minimum of 200 Units of AfIII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

tephonie Onetto

Stephanie Cornelio Production Scientist 20 Jun 2018

M YILan

Mary Conton Packaging Quality Control Inspector 20 Jun 2018

