

## New England Biolabs Certificate of Analysis

**Product Name:** BsgI  
**Catalog Number:** R0559S  
**Concentration:** 5,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total volume of 50 µl.  
**Packaging Lot Number:** 10098888  
**Expiration Date:** 02/2023  
**Storage Temperature:** -20°C  
**Storage Conditions:** 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 0.32 mM S-adenosylmethionine (SAM), 50% Glycerol, 500 µg/ml BSA (pH 7.4 @ 25°C)  
**Specification Version:** PS-R0559S/L v2.0

BsgI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0559SVIAL	BsgI	10098887	Pass
B7204SVIAL	CutSmart® Buffer	10093127	Pass

Assay Name/Specification	Lot # 10098888
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 50 units of BsgI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 5-fold over-digestion of Lambda DNA with BsgI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with BsgI.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 15 Units of BsgI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Pengda Zhang  
Production Scientist  
01 Mar 2021



Michael Tonello  
Packaging Quality Control Inspector  
01 Mar 2021