

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | Nt.BbvCl   |
|------------------------|--|
| Catalog Number:        | R0632L   |
| Concentration:         | 10,000 U/ml  |
| Unit Definition:       | One unit is defined as the amount of enzyme required to convert 1 μg<br>of supercoiled pUB DNA to open circular form in 1 hour at 37°C in a<br>total reaction volume of 50 μl. |
| Packaging Lot Number:  | 10238162   |
| Expiration Date:       | 02/2026  |
| Storage Temperature:   | -20°C  |
| Storage Conditions:    | 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%<br>Glycerol, 200 μg/ml BSA  |
| Specification Version: | PS-R0632S/L v1.0   |

| Nt.BbvCl Component List |                       |            |                      |
|-------------------------|-----------------------|------------|----------------------|
| NEB Part Number         | Component Description | Lot Number | Individual QC Result |
| R0632LVIAL              | Nt.BbvCl              | 10226116   | Pass                 |
| B6004SVIAL              | rCutSmart™ Buffer     | 10233336   | Pass                 |

| Assay Name/Specification  | Lot # 10238162 |
|---|----------------|
| Endonuclease Activity (Nicking)<br>A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of supercoiled pUC19 DNA and a<br>minimum of 30 units of Nt.BbvCl incubated for 4 hours at 37°C results in <10%<br>conversion to the nicked form as determined by agarose gel electrophoresis.                | Pass           |
| Exonuclease Activity (Radioactivity Release)<br>A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of a mixture of single and<br>double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 30 units of Nt.BbvCI incubated<br>for 4 hours at 37°C releases <0.1% of the total radioactivity.    | Pass           |
| Non-Specific DNase Activity (16 Hour)<br>A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of Lambda DNA and a minimum of<br>30 Units of Nt.BbvCl incubated for 16 hours at 37°C results in a DNA pattern free of<br>detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.





240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

11.

Ana Egana Production Scientist 28 Mar 2024

Michae 111

Michael Tonello Packaging Quality Control Inspector 28 Mar 2024

