

## New England Biolabs Certificate of Analysis

**Product Name:** Sall-HF<sup>®</sup>  
**Catalog Number:** R3138S  
**Concentration:** 20,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (HindIII digest) in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10193702  
**Expiration Date:** 04/2025  
**Storage Temperature:** -20°C  
**Storage Conditions:** 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 300 µg/ml BSA, (pH 7.5 @ 25°C)  
**Specification Version:** PS-R3138S/L/V v2.0


Sall-HF <sup>®</sup> Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R3138SVIAL	Sall-HF <sup>®</sup>	10183138	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10189226	Pass
B6004SVIAL	rCutSmart <sup>™</sup> Buffer	10193042	Pass

Assay Name/Specification	Lot # 10193702
<b>Blue-White Screening (Terminal Integrity)</b> A sample of pUC19 vector linearized with a 10-fold excess of Sall-HF <sup>®</sup> , religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in CutSmart <sup>®</sup> Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of Sall-HF <sup>®</sup> incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart <sup>®</sup> Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H ] E. coli DNA and a minimum of 200 units of Sall-HF <sup>®</sup> incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 50-fold over-digestion of pBC4XS DNA with Sall-HF <sup>®</sup> , >95% of the DNA	Pass

Assay Name/Specification	Lot # 10193702
<p>fragments can be ligated with T4 DNA ligase in 4 hours at 25°C. Of these ligated fragments, &gt;95% can be recut with Sall-HF®.</p> <p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of pBR322 DNA and a minimum of 200 units of Sall-HF® incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



YunJie Sun  
Production Scientist  
12 Apr 2023



Michael Tonello  
Packaging Quality Control Inspector  
18 Jul 2023