

New England Biolabs Product Specification

Product Name: NEB[®] 10-beta Electrocompetent *E. coli*
Catalog #: C3020K
Shelf Life: 12 months
Storage Temp: -80°C
Specification Version: PS-C3020K v1.0
Effective Date: 17 Jan 2017

Assay Name/Specification (minimum release criteria)

Antibiotic Resistance (Streptomycin) - 15 µl of untransformed NEB[®] 10-beta Electrocompetent *E. coli* streaked onto a LB or Rich Broth plate containing Streptomycin will form colonies after incubation for 16 hours at 37°C.

Antibiotic Sensitivity (Ampicillin) - 15 µl of untransformed NEB[®] 10-beta Electrocompetent *E. coli* streaked onto a LB or Rich Broth plate containing Ampicillin will not form colonies after incubation for 16 hours at 37°C.

Antibiotic Sensitivity (Chloramphenicol) - 15 µl of untransformed NEB[®] 10-beta Electrocompetent *E. coli* streaked onto a LB or Rich Broth plate containing Chloramphenicol will not form colonies after incubation for 16 hours at 37°C.

Antibiotic Sensitivity (Kanamycin) - 15 µl of untransformed NEB[®] 10-beta Electrocompetent *E. coli* streaked onto a LB or Rich Broth plate containing Kanamycin will not form colonies after incubation for 16 hours at 37°C.

Antibiotic Sensitivity (Nitrofurantoin) - 15 µl of untransformed NEB[®] 10-beta Electrocompetent *E. coli* streaked onto a LB or Rich Broth plate containing Nitrofurantoin will not form colonies after incubation for 16 hours at 37°C.

Antibiotic Sensitivity (Spectinomycin) - 15 µl of untransformed NEB[®] 10-beta Electrocompetent *E. coli* streaked onto a LB or Rich Broth plate containing Spectinomycin will not form colonies after incubation for 16 hours at 37°C.

Antibiotic Sensitivity (Tetracycline) - 15 µl of untransformed NEB[®] 10-beta Electrocompetent *E. coli* streaked onto a LB or Rich Broth plate containing Tetracycline will not form colonies after incubation for 16 hours at 37°C.

Blue-White Screening (α-complementation, Competent Cells) - NEB[®] 10-beta Electrocompetent *E. coli* were shown to be suitable for blue/white screening by α-complementation of the β-galactosidase gene using pUC19.

Phage Resistance (Φ 80) - 15 µl of untransformed NEB[®] 10-beta Electrocompetent *E. coli* streaked onto a Rich Broth plate does not support plaque formation by phage Φ 80 after incubation for 16 hours at 37°C.

Transformation Efficiency - 25 µl of NEB[®] 10-beta Electrocompetent *E. coli* cells were transformed with 10 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB-Ampicillin plates at 37°C resulted in >2 x 10¹⁰ cfu/µg of DNA.



Date 17 Jan 2017

Derek Robinson
Director of Quality Control

