NEW ENGLAND Biolabs®

240 County Road
Ipswich, MA 01938-2723

Tel 978-927-5054
Fax 978-921-1350

www.neb.com
info@neb.com

New England Biolabs
Product Specification

Product Name:	*BsaBI*
Catalog #:	*R0537S/L/V*
Concentration:	*10,000 units/ml*
Unit Definition:	*One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (dam-) in 1 hour at 60°C in a total reaction volume of 50 µl.*
Shelf Life:	*24 months*
Storage Temp:	*-20°C*
Storage Conditions:	*300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml BSA*
Specification Version:	*PS-R0537S/L v1.0*
Effective Date:	*06/11/2013*

Assay Name/Specification (minimum release criteria)
Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [^{3}H] *E. coli* DNA and a minimum of 100 units of BsaBI incubated for 4 hours at 60ºC releases <0.1% of the total radioactivity.
Ligation and Recutting (Terminal Integrity) - After a 10-fold over-digestion of Lambda dam- DNA with BsaBI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16ºC. Of these ligated fragments, >95% can be recut with BsaBI.
Non-Specific DNase Activity (16 hour) - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda dam- DNA and a minimum of 10 units of BsaBI incubated for 16 hours at 60ºC results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.
Protein Purity Assay (SDS-PAGE) - BsaBI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

Date 06/11/2013

Derek Robinson
Quality Approver