240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Product Specification

Product Name: SrfI

Catalog #: R0629S/L
Concentration: 20,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of pNEB193-SrfI DNA in CutSmart incubated for 1 hour

at 37°C in a total reaction volume of 50  $\mu$ l.

Shelf Life: 18 months
Storage Temp: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 500 µg/ml BSA, (pH 7.4 @ 25°C)

Specification Version: PS-R0629S/L v1.0

Effective Date: 11 Nov 2015

## Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50  $\mu$ l reaction in CutSmart® Buffer containing 1  $\mu$ g of supercoiled pBR322 DNA and a minimum of 100 units of Srfl incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup>H] *E. coli* DNA and a minimum of 200 units of Srfl incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Functional Testing (15 minute Digest) - A 50 μl reaction in CutSmart® Buffer containing 1 μg of pNEB193-SrfI DNA and 1 μl of SrfI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.

**Ligation and Recutting (Terminal Integrity)** - After a 20-fold over-digestion of pNEB193-SrfI DNA with SrfI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with SrfI.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmart® Buffer containing 1 µg of pNEB193-SrfI DNA and a minimum of 20 units of SrfI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Date

11 Nov 2015

Protein Purity Assay (SDS-PAGE) - Srfl is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

Derek Robinson

Director of Quality Control





