

## New England Biolabs Product Specification

**Product Name:** *I-CeuI*  
**Catalog #:** *R0699S/L*  
**Concentration:** *5,000 units/ml*  
**Unit Definition:** *One unit is defined as the amount of enzyme required to cleave 1 µg of pBHS ScaI-linearized Control Plasmid in 3 hours at 37°C in a total reaction volume of 50 µl.*  
**Shelf Life:** *24 months*  
**Storage Temp:** *-20°C*  
**Storage Conditions:** *300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml BSA*  
**Specification Version:** *PS-R0699S/L v1.0*  
**Effective Date:** *07/29/2013*

### Assay Name/Specification (minimum release criteria)

**Endonuclease Activity (Nicking)** - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 15 Units of I-CeuI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.

**Exonuclease Activity (Radioactivity Release)** - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA and a minimum of 50 units of I-CeuI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

**Ligation and Recutting (Terminal Integrity)** - After a 10-fold over-digestion of pBHS-ScaI DNA with I-CeuI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with I-CeuI.

**Non-Specific DNase Activity (16 Hour)** - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBHS-ScaI DNA and a minimum of 50 Units of I-CeuI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

One or more products referenced in this document may be covered by a 3rd-party trademark.  
Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Date 07/29/2013

Derek Robinson  
Quality Approver

