NEW ENGLAND BioLabs® Inc.

240 County Road
Ipswich, MA 01938-2723

Tel 978-927-5054
Fax 978-921-1350

www.neb.com
info@neb.com

New England Biolabs
Product Specification

Product Name:	*BmtI-HF®*
Catalog #:	*R3658S/L*
Concentration:	*20,000 units/ml*
Unit Definition:	*One unit is defined as the amount of enzyme required to digest 1 µg of pXba in 1 hour at 37°C in a total reaction volume of 50 µl.*
Shelf Life:	*24 months*
Storage Temp:	*-20 °C*
Storage Conditions:	*300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml BSA*
Specification Version:	*PS-R3658S/L v1.0*
Effective Date:	*03 May 2013*

Assay Name/Specification (minimum release criteria)

Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [^{3}H] *E. coli* DNA and a minimum of 200 units of BmtI-HF™ incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Ligation and Recutting (Terminal Integrity) - After a 20-fold over-digestion of pXba DNA with BmtI-HF™, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BmtI-HF™.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of 100 Units of BmtI-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

* *The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.*

Date 03 May 2013

Derek Robinson
Director of Quality Control