

## New England Biolabs Certificate of Analysis

**Product Name:** NEBuffer™ 2  
**Catalog Number:** B7002S  
**Concentration:** 10 X Concentrate  
**Packaging Lot Number:** 10105382  
**Expiration Date:** 02/2024  
**Storage Temperature:** -20°C  
**Specification Version:** PS-B7002S v1.0  
**Composition (1X):** 50 mM NaCl, 10 mM Tris-HCl, 10 mM MgCl<sub>2</sub>, 1 mM DTT, (pH 7.9 @ 25°C)

NEBuffer™ 2 Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
B7002SVIAL	NEBuffer™ 2	10097265	Pass

Assay Name/Specification	Lot # 10105382
<b>RNase Activity (Buffer)</b> A 10 µl reaction in 1X NEBuffer 2 containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection.	Pass
<b>Non-Specific DNase Activity (16 hour, Buffer)</b> A 50 µl reaction in 1X NEBuffer 2 containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>pH (buffers/solutions)</b> The pH of 10X NEBuffer 2 is between pH 7.8 and 8.0 at 25°C.	Pass
<b>Endonuclease Activity (Nicking, Buffer)</b> A 50 µl reaction in 1X NEBuffer 2 containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Conductivity (buffers/solutions)</b> The conductivity of 10X NEBuffer 2 is between 52 and 78 mS/cm at 25°C.	Pass
<b>Functional Testing (Restriction Digest, BSA, Buffer)</b> A 50 µl reaction in 1X NEBuffer 2 plus 100 µg/ml Bovine Serum Albumin containing 1	Pass

Assay Name/Specification	Lot # 10105382
<p>µg of Lambda DNA and 1 unit of HindIII incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.</p> <p><b>Functional Testing (Restriction Digest, BSA, Buffer)</b> A 50 µl reaction in 1X NEBuffer 2 plus 100 µg/ml Bovine Serum Albumin containing 1 µg of Lambda DNA and 1 unit of SphI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Michael Dalton  
Production Scientist  
29 Apr 2021



Michael Tonello  
Packaging Quality Control Inspector  
29 Apr 2021