

## New England Biolabs Certificate of Analysis


**Product Name:** Exonuclease III (*E.coli*)  
**Catalog Number:** M0206L  
**Concentration:** 100,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to produce 1 nmol of acid-soluble total nucleotide in a total reaction volume of 50 µl in 30 minutes at 37°C in 1X NEBuffer 1 with 0.15 mM sonicated duplex [ 3H]-DNA.  
**Packaging Lot Number:** 10085452  
**Expiration Date:** 02/2022  
**Storage Temperature:** -20°C  
**Storage Conditions:** 200 mM KCl, 5 mM KPO<sub>4</sub>, 0.05 mM EDTA, 5 mM βME, 50 % Glycerol, 200 µg/ml BSA, (pH 6.5 @ 25°C)  
**Specification Version:** PS-M0206S/L v1.0

| Exonuclease III ( <i>E.coli</i> ) Component List |                                   |            |                      |
|--|-----------------------------------|------------|----------------------|
| NEB Part Number                                  | Component Description             | Lot Number | Individual QC Result |
| M0206LVIAL                                       | Exonuclease III ( <i>E.coli</i> ) | 10063239   | Pass                 |
| B7001SVIAL                                       | NEBuffer™ 1                       | 10064408   | Pass                 |

| Assay Name/Specification  | Lot # 10085452 |
|---|----------------|
| <p><b>qPCR DNA Contamination (<i>E. coli</i> Genomic)</b><br/>           A minimum of 100 units of Exonuclease III (<i>E. coli</i>) is screened for the presence of <i>E. coli</i> genomic DNA using SYBR® Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is ≤ 1 <i>E. coli</i> genome.</p> | Pass           |
| <p><b>Protein Purity Assay (SDS-PAGE)</b><br/>           Exonuclease III (<i>E. coli</i>) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>  | Pass           |
| <p><b>Endonuclease Activity (Nicking)</b><br/>           A 50 µl reaction in NEBuffer 1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 300 units of Exonuclease III (<i>E. coli</i>) incubated for 4 hours at 37°C results in &lt;20% conversion to the nicked form as determined by agarose gel electrophoresis.</p>  | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



---

John Greci  
Production Scientist  
08 Oct 2020



---

Michael Tonello  
Packaging Quality Control Inspector  
08 Oct 2020