

## New England Biolabs Certificate of Analysis

**Product Name:** DNA Polymerase I (*E. coli*)  
**Catalog #:** M0209S/L  
**Concentration:** 10,000 units/ml  
**Unit Definition:** One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 37°C.  
**Lot #:** 0991703  
**Assay Date:** 03/2017  
**Expiration Date:** 3/2019  
**Storage Temp:** -20°C  
**Storage Conditions:** 25 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, (pH 7.4 @ 25°C)  
**Specification Version:** PS-M0209S/L v1.0  
**Effective Date:** 02 May 2016

Assay Name/Specification (minimum release criteria)	Lot #0991703
<b>Endonuclease Activity (Nicking)</b> - A 50 µl reaction in NEBuffer 2 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 units of DNA Polymerase I ( <i>E. coli</i> ) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Phosphatase Activity (pNPP)</b> - A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl <sub>2</sub> containing 2.5 mM <i>p</i> -Nitrophenyl Phosphate (pNPP) and a minimum of 100 units DNA Polymerase I ( <i>E. coli</i> ) incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	<b>Pass</b>
<b>Protein Purity Assay (SDS-PAGE)</b> - DNA Polymerase I ( <i>E. coli</i> ) is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	<b>Pass</b>
<b>qPCR DNA Contamination (<i>E. coli</i> Genomic)</b> - A minimum of 10 units of DNA Polymerase I ( <i>E. coli</i> ) is screened for the presence of <i>E. coli</i> genomic DNA using SYBR® Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is ≤ 1 <i>E. coli</i> genome.	<b>Pass</b>
<b>RNase Activity (Extended Digestion)</b> - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of DNA Polymerase I ( <i>E. coli</i> ) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	<b>Pass</b>



Authorized by  
Melanie Fortier  
02 May 2016



Inspected by  
Tony Spear-Alfonso  
28 Mar 2017

