

New England Biolabs Certificate of Analysis

Product Name: Antarctic Phosphatase
Catalog Number: M0289S
Concentration: 5,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme that will dephosphorylate 1 µg of pUC19 vector DNA cut with a restriction enzyme generating 5' recessed ends in 30 minutes at 37°C. Dephosphorylation is defined as >95% inhibition of recircularization in a self-ligation reaction and is measured by transformation into *E. coli*.
Packaging Lot Number: 10144347
Expiration Date: 03/2024
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl, 1 mM MgCl₂, 0.01 mM ZnCl₂, 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version: PS-M0289S/L v2.0

Antarctic Phosphatase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0289SVIAL	Antarctic Phosphatase	10140300	Pass
B0289SVIAL	Antarctic Phosphatase Reaction Buffer	10140303	Pass

Assay Name/Specification	Lot # 10144347
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Antarctic Phosphatase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 4 containing 1 µg of PhiX174-HaeIII DNA and a minimum of 50 units of Antarctic Phosphatase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of Antarctic Phosphatase incubated for 4 hours at 37°C results	Pass

Assay Name/Specification	Lot # 10144347
in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
<p>Protein Purity Assay (SDS-PAGE) Antarctic Phosphatase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in Antarctic Phosphatase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of Antarctic Phosphatase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 5 units of Antarctic Phosphatase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Ana Egana
Production Scientist
08 Apr 2022



Josh Hersey
Packaging Quality Control Inspector
08 Apr 2022