

## New England Biolabs Certificate of Analysis

**Product Name:** Bst 2.0 DNA Polymerase  
**Catalog Number:** M0537L  
**Concentration:** 8,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme that will incorporate 25 nmol of dNTP into acid insoluble material in 30 minutes at 65°C.  
**Packaging Lot Number:** 10131474  
**Expiration Date:** 09/2023  
**Storage Temperature:** -20°C  
**Storage Conditions:** 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.1 % Triton®X-100, 50 % Glycerol, (pH 7.1 @ 25°C)  
**Specification Version:** PS-M0537S/L v2.0

Bst 2.0 DNA Polymerase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0537LVIAL	Bst 2.0 DNA Polymerase	10128902	Pass
B1003SVIAL	Magnesium Sulfate (MgSO <sub>4</sub> ) Solution	10118450	Pass
B0537SVIAL	Isothermal Amplification Buffer	10128740	Pass

Assay Name/Specification	Lot # 10131474
<p><b>Exonuclease Activity (Radioactivity Release)</b>            A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 500 units of Bst 2.0 DNA Polymerase incubated for 4 hours at 65°C releases &lt;0.1% of the total radioactivity.</p>	Pass
<p><b>Endonuclease Activity (Nicking)</b>            A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 500 units of Bst 2.0 DNA Polymerase incubated for 4 hours at 65°C results in &lt;10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p><b>RNase Activity (Extended Digestion)</b>            A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Bst 2.0 DNA Polymerase is incubated at 37°C. After incubation for 16 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass


Assay Name/Specification	Lot # 10131474
<p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 120 units of Bst 2.0 DNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Protein Purity Assay (SDS-PAGE)</b> Bst 2.0 DNA Polymerase is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>Phosphatase Activity (pNPP)</b> A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl<sub>2</sub> containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units Bst 2.0 DNA Polymerase incubated for 4 hours at 37°C yields &lt;0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	<b>Pass</b>
<p><b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 120 units of Bst 2.0 DNA Polymerase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Christie Vazquez  
Production Scientist  
29 Nov 2021



Michael Tonello  
Packaging Quality Control Inspector  
29 Nov 2021