

## New England Biolabs Certificate of Analysis

**Product Name:** Streptavidin  
**Catalog Number:** N7021S  
**Concentration:** 1 mg/ml  
**Packaging Lot Number:** 10115460  
**Expiration Date:** 07/2023  
**Storage Temperature:** -20°C  
**Storage Conditions:** 140 mM NaCl, 8 mM Sodium Phosphate, 2 mM Potassium Phosphate, 10 mM KCl, (pH 7.4 @ 25°C)  
**Specification Version:** PS-N7021S v2.0

Streptavidin Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
N7021SVIAL	Streptavidin	10115459	Pass

Assay Name/Specification	Lot # 10115460
<b>Specific Activity</b> 1 mg of Streptavidin is required to bind $\geq 14$ $\mu\text{g}$ of Biotin.	Pass
<b>Protein Purity Assay (SDS-PAGE)</b> Streptavidin is $\geq 95\%$ pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 $\mu\text{l}$ reaction in NEBuffer 3 containing 1 $\mu\text{g}$ of Lambda DNA and a minimum of 1 $\mu\text{g}$ of Streptavidin incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>RNase Activity (Extended Digestion)</b> A 10 $\mu\text{l}$ reaction in NEBuffer 3 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 $\mu\text{g}$ of Streptavidin is incubated at 37°C. After incubation for 2 hours, $>90\%$ of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
<b>Functional Testing (Single Stranded DNA Binding - FAM Labeled Oligo)</b> A 20 $\mu\text{l}$ reaction in NEBuffer 3 containing 3 $\mu\text{M}$ FAM and Biotin-labeled 50-mer and a maximum of 1 $\mu\text{g}$ of Streptavidin incubated for 5 minutes at 25°C produces a mobility shift in $>95\%$ of the starting material as determined by TBE gel electrophoresis and	Pass

Assay Name/Specification	Lot # 10115460
<p>UV imaging.</p> <p><b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer 3 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 1 µg of Streptavidin incubated for 4 hours at 37°C results in &lt;10% conversion to the nicked form as determined by agarose gel electrophoresis.</p> <p><b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in NEBuffer 3 containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 1 µg of Streptavidin incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p>	<p style="text-align: center;"><b>Pass</b></p> <p style="text-align: center;"><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Bo Wu  
Production Scientist  
04 Aug 2021



Michael Tonello  
Packaging Quality Control Inspector  
04 Aug 2021