

New England Biolabs Certificate of Analysis

Product Name: α -N-Acetylgalactosaminidase
Catalog #: P0734S/L
Concentration: 20,000 units/ml
Unit Definition: One unit is defined as the amount of enzyme required to cleave > 95% of the terminal α -D-N-acetylgalactosamine from 1 nmol (GalNAc α 1-3)(Fuc α 1-2)Gal β 1-4Glc-7-amino-4-methyl-coumarin (AMC), in 1 hour at 37°C in a total reaction volume of 10 μ l.
Lot #: 0071703
Assay Date: 03/2017
Expiration Date: 3/2019
Storage Temp: -20°C
Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, (pH 7.5 @ 25°C)
Specification Version: PS-P0734S/L v1.0
Effective Date: 17 Feb 2016

Assay Name/Specification (minimum release criteria)	Lot #0071703
Glycosidase Activity (Endo F1, F2, H) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 20 units of α -N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (Endo F2, F3) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 20 units of α -N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (PNGase F) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled PNGase F substrate (Fluoresceinated fetuin triantennary) and 20 units of α -N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-Mannosidase) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β -Mannosidase substrate (Man β 1-4Man β 1-4Man-AMC) and 20 units of α -N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-Xylosidase) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β -Xylosidase substrate (Xyl β 1-4Xyl β 1-4Xyl β 1-4Xyl-AMC) and 20 units of α -N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass



New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0071703
Glycosidase Activity (β1-3 Galactosidase) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β -Galactosidase substrate (Gal β 1-3GlcNAc β 1-4Gal β 1-4Glc-AMC) and 20 units of α -N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-4 Galactosidase) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β -Galactosidase substrate (Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc-AMC) and 20 units of α -N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-N-Acetylgalactosaminidase) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β -N-Acetylgalactosaminidase substrate (GalNAc β 1-4Gal β 1-4Glc-AMC) and 20 units of α -N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-N-Acetylglucosaminidase) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β -N-Acetylglucosaminidase substrate (GlcNAc β 1-4GlcNAc β 1-4GlcNAc-AMC) and 20 units of α -N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α-Glucosidase) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α -Glucosidase substrate (Glc α 1-6Glc α 1-4Glc-AMC) and 20 units of α -N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α-Neuraminidase) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α -Neuraminidase substrate (Neu5Ac α 2-3Gal β 1-3GlcNAc β 1-3Gal β 1-4Glc-AMC) and 20 units of α -N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-2 Fucosidase) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α -Fucosidase substrate (Fuc α 1-2Gal β 1-4Glc-AMC) and 20 units of α -N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-3 Fucosidase) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α -Fucosidase substrate (Fuc α 1-3Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc-AMC) and 20 units of α -N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass

New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0071703
<p>Glycosidase Activity (α1-3 Galactosidase) - A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-3Galβ1-4GlcNAc-AMC) and 20 units of α-N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Mannosidase) - A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-3Manβ1-4GlcNAc-AMC) and 20 units of α-N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-6 Galactosidase) - A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-6Galα1-6Glcα1-2Fru-AMC) and 20 units of α-N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-6 Mannosidase) - A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-6Manα1-6(Manα1-3)Man-AMC) and 20 units of α-N-Acetylgalactosaminidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Protease Activity (SDS-PAGE) - A 20 μl reaction in 1X Glyco Buffer 1 containing 24 μg of a standard mixture of proteins and a minimum of 100 units of α-N-Acetylgalactosaminidase incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) - α-N-Acetylgalactosaminidase is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass



Authorized by
Derek Robinson
17 Feb 2016



Inspected by
Alicia Bielik
24 Mar 2017

