

## New England Biolabs Certificate of Analysis

**Product Name:** *FspI*  
**Catalog #:** R0135S/L  
**Concentration:** 10,000 units/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Lot #:** 0601405  
**Assay Date:** 05/2014  
**Expiration Date:** 05/2016  
**Storage Temp:** -20 °C  
**Storage Conditions:** 300mM NaCl, 10mM Tris-HCl (pH 7.5), 0.1mM EDTA, 1mM dithiothreitol, 0.15% Triton X-100, 300 ug/ml BSA, 50% glycerol  
**Specification Version:** PS-R0135S/L v1.0  
**Effective Date:** 08 Jul 2013

Assay Name/Specification (minimum release criteria)	Lot #0601405
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 100 units of FspI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	<b>Pass</b>
<b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of Lambda DNA with FspI, ~50% of the DNA fragments can be ligated with T4 DNA ligase in 4 hours at 25°C. Of these ligated fragments, >95% can be recut with FspI.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 Hour)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 Units of FspI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	<b>Pass</b>

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



Authorized by  
Derek Robinson  
08 Jul 2013



Inspected by  
David Hough  
05 Sep 2014

