

New England Biolabs Certificate of Analysis

Product Name: KpnI
Catalog Number: R0142S
Concentration: 10,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10107517
Expiration Date: 04/2023
Storage Temperature: -20°C
Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA
Specification Version: PS-R0142S/L v2.0

KpnI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0142SVIAL	KpnI	10107516	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10093116	Pass
B6001SVIAL	NEBuffer™ r1.1	10102943	Pass

Assay Name/Specification	Lot # 10107517
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pXba DNA with KpnI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with KpnI.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of pXba DNA and a minimum of 50 Units of KpnI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) KpnI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
Blue-White Screening (Terminal Integrity) A sample of pUC19 vector linearized with a 10-fold excess of KpnI, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass

Assay Name/Specification	Lot # 10107517
<p>Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 units of KpnI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of KpnI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Penghua Zhang
Production Scientist
17 May 2021



Michael Tonello
Packaging Quality Control Inspector
17 May 2021