

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Bglll
Catalog Number: R0144S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10065427
Expiration Date: 08/2021
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0144S/L v2.0

BgIII Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0144SVIAL	BgIII	10053466	Pass	
B7203SVIAL	NEBuffer™ 3.1	10053974	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10065745	Pass	

Assay Name/Specification	Lot # 10065427
Protein Purity Assay (SDS-PAGE) Bglll is >95% pure as determined by SDS PAGE analysis using Coomassie Blue	Pass
detection.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 100 Units of BgIII incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 20-fold over-digestion of Lambda DNA with BgIII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
>95% can be recut with BgIII.	
Blue-White Screening (Terminal Integrity)	Pass
A sample of LITMUS28i vector linearized with a 10-fold excess of BgIII, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene	
results in <1% white colonies.	



R0144S / Lot: 10065427

Page 1 of 2

Assay Name/Specification	Lot # 10065427
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 3.1 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 10 Units of BglII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in NEBuffer 3.1 containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of BgIII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass

This product has been tested and shown to be in compliance with all specifications.

Penghaa Zhang Production Scientist

28 Aug 2019

Michael Tonello

Packaging Quality Control Inspector

18 Mar 2020



R0144S / Lot: 10065427

Page 2 of 2