

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Clal
Catalog Number: R0197S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 μg

of Lambda DNA (dam-) in 1 hour at 37°C in a total reaction volume of

50 μl.

Packaging Lot Number: 10117183
Expiration Date: 04/2023
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0197S/L v1.0

Clal Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0197SVIAL	Clal	10104969	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10108731	Pass	
B6004SVIAL	rCutSmart™ Buffer	10114153	Pass	

Assay Name/Specification	Lot # 10117183
Protein Purity Assay (SDS-PAGE)	Pass
Clal is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	
a minimum of 30 Units of Clal incubated for 4 hours at 37°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	
double-stranded [³H] E. coli DNA and a minimum of 100 units of Clal incubated for 4	
hours at 37°C releases <0.1% of the total radioactivity.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 μl reaction in CutSmart™ Buffer containing 1 μg of Lambda dam- DNA and a	
minimum of 100 Units of Clal incubated for 16 hours at 37°C results in a DNA pattern	
free of detectable nuclease degradation as determined by agarose gel	



R0197S / Lot: 10117183

Page 1 of 2

Assay Name/Specification	Lot # 10117183
electrophoresis.	
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda dam- DNA with Clal, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Clal.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

16 Aug 2021

Michael Tonello

Packaging Quality Control Inspector

16 Aug 2021



R0197S / Lot: 10117183

Page 2 of 2