

New England Biolabs Certificate of Analysis

Product Name: PpuMI
Catalog Number: R0506S
Concentration: 10,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (HindIII digest) in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10236222
Expiration Date: 03/2026
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl, 50 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)
Specification Version: PS-R0506S/L v2.0


| PpuMI Component List | | | |
|----------------------|-----------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| R0506SVIAL | PpuMI | 10230750 | Pass |
| B6004SVIAL | rCutSmart™ Buffer | 10229453 | Pass |

| Assay Name/Specification | Lot # 10236222 |
|--|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 units of PpuMI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of PpuMI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Functional Testing (15 minute Digest) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda-HindIII DNA and 1 µl of PpuMI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pBC4 DNA with PpuMI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% | Pass |

| Assay Name/Specification | Lot # 10236222 |
|---|--------------------|
| <p>can be recut with PpuMI.</p> <p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 100 units of PpuMI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p> | <p>Pass</p> |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Ana Egana
Production Scientist
11 Mar 2024



Michael Tonello
Packaging Quality Control Inspector
11 Mar 2024