

## New England Biolabs Certificate of Analysis

**Product Name:** *ApaLI*  
**Catalog Number:** *R0507M*  
**Concentration:** *50,000 U/ml*  
**Unit Definition:** *One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (HindIII digest) in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.*  
**Packaging Lot Number:** *10149249*  
**Expiration Date:** *05/2024*  
**Storage Temperature:** *-20°C*  
**Storage Conditions:** *10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)*  
**Specification Version:** *PS-R0507T/M v2.0*

ApaLI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0507MVIAL	ApaLI	10149247	Pass
B6004SVIAL	rCutSmart™ Buffer	10143288	Pass

Assay Name/Specification	Lot # 10149249
<p><b>Ligation and Recutting (Terminal Integrity)</b>            After a 10-fold over-digestion of Lambda DNA with ApaLI, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with ApaLI.</p>	Pass
<p><b>Endonuclease Activity (Nicking)</b>            A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled M13mp18 DNA and a minimum of 50 units of ApaLI incubated for 4 hours at 37°C results in &lt;10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p><b>Exonuclease Activity (Radioactivity Release)</b>            A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 100 units of ApaLI incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p>	Pass
<p><b>Functional Testing (15 minute Digest)</b>            A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda-HindIII DNA and 1 µl of ApaLI incubated for 15 minutes at 37°C results in complete digestion as</p>	Pass

Assay Name/Specification	Lot # 10149249
<p>determined by agarose gel electrophoresis.</p> <p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 100 units of ApaLI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.




---

Penghua Zhang  
Production Scientist  
06 Jun 2022




---

Michael Tonello  
Packaging Quality Control Inspector  
06 Jun 2022