

New England Biolabs Certificate of Analysis

Product Name: *BstBI*
Catalog Number: *R0519L*
Concentration: *20,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in rCutSmart Buffer in 1 hour at 65°C in a total reaction volume of 50 µl.*
Packaging Lot Number: *10206902*
Expiration Date: *09/2025*
Storage Temperature: *-20°C*
Storage Conditions: *10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)*
Specification Version: *PS-R0519S/L v2.0*

BstBI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0519LVIAL	BstBI	10206315	Pass
B6004SVIAL	rCutSmart™ Buffer	10202500	Pass

Assay Name/Specification	Lot # 10206902
<p>Endonuclease Activity (Nicking) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled pUC19 DNA and a minimum of 20 units of BstBI incubated for 4 hours at 65°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of BstBI incubated for 4 hours at 65°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Functional Testing (15 minute Digest) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 µl of BstBI incubated for 15 minutes at 65°C results in complete digestion as determined by agarose gel electrophoresis.</p>	Pass
<p>Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with BstBI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,</p>	Pass

Assay Name/Specification	Lot # 10206902
<p>>95% can be recut with BstBI.</p> <p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 20 units of BstBI incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



YunJie Sun
Production Scientist
11 Sep 2023



Josh Hersey
Packaging Quality Control Inspector
29 Sep 2023