

New England Biolabs Certificate of Analysis

Product Name: BsiWI
Catalog #: R0553S/L
Concentration: 10,000 units/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of PhiX174 DNA in 1 hour at 55°C in a total reaction volume of 50 µl.
Lot #: 0341701
Assay Date: 01/2017
Expiration Date: 1/2019
Storage Temp: -20°C
Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml BSA
Specification Version: PS-R0553S/L v1.0
Effective Date: 28 Oct 2014

Assay Name/Specification (minimum release criteria)	Lot #0341701
Endonuclease Activity (Nicking) - A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled pUC19 DNA and a minimum of 10 Units of BsiWI incubated for 4 hours at 55°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 20 units of BsiWI incubated for 4 hours at 55°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) - After a 10-fold over-digestion of PhiX174 DNA with BsiWI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BsiWI.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 3.1 containing 1 µg of PhiX174 DNA and a minimum of 10 Units of BsiWI incubated for 16 hours at 55°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass



Authorized by
Derek Robinson
28 Oct 2014



Inspected by
Jianying Luo
10 Feb 2017

