

drive DISCOVERY

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## **New England Biolabs Certificate of Analysis**

Product Name: SgrAI Catalog Number: R0603L 10,000 U/ml Concentration:

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of LambdaDNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10138028 Expiration Date: 10/2023 -20°C Storage Temperature:

100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Storage Conditions:

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0603S/L v1.0

SgrAl Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0603LVIAL	SgrAI	10122894	Pass	
B6004SVIAL	rCutSmart™ Buffer	10132778	Pass	

Assay Name/Specification	Lot # 10138028
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 50 units of SgrAl incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 30 Units of SgrAl incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 2-fold over-digestion of Lambda DNA with SgrAI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with SgrAI.	Pass
Non-Specific DNase Activity (16 hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 10 Units of SgrAl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE:	Pass



R0603L / Lot: 10138028

Page 1 of 2



Assay Name/Specification	Lot # 10138028
although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity.	
See the product FAQ for recommended reaction conditions for this enzyme.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

17 Feb 2022

Michael Tonello

Packaging Quality Control Inspector

17 Feb 2022